


Innate Immunity

Moumita Pal

Immune system is one of the most crucial systems of the body that is very important for survival. It encompasses tissues and cells that are associated with the defense of the body from different pathogens and infectious agents. The immunity is generally classified into two different types: namely, **innate immunity and adaptive immunity**.



Innate Immunity

Innate immunity is also referred to as immediate response immune system. This system activates within minutes to hours after a foreign agent invades within the body. Innate immune system is composed of two lines of defenses. First line of defense consists of skin (mainly the epidermis), gastric acid in the stomach and mucus membranes. The second line of defense consists of chemicals and cells that are released in the blood after being exposed to a pathogenic stimulus.

Innate Immune System


The innate immune system is made of defenses against infection that can be activated immediately once a pathogen attacks. The innate immune system is essentially made up of barriers that aim to keep viruses, bacteria, parasites, and other foreign particles out of your body or limit their ability to spread and move throughout the body. The innate immune system includes:

- *Physical Barriers*
 - such as skin, the gastrointestinal tract, the respiratory tract, the nasopharynx, cilia, eyelashes and other body hair.
- *Defense Mechanisms*
 - such as secretions, mucous, bile, gastric acid, saliva, tears, and sweat.
- *General Immune Responses*
 - such as inflammation, complement, and non-specific cellular responses. The inflammatory response actively brings immune cells to the site of an infection by increasing blood flow to the area. Complement is an immune response that marks pathogens for destruction and makes holes in the cell membrane of the pathogen.


The innate immune system is always general, or *nonspecific*, meaning anything that is identified as foreign or *non-self* is a target for the innate immune response. The innate immune system is activated by the presence of antigens and their chemical properties.

Cells of the Innate Immune System


There are many types of white blood cells, or *leukocytes*, that work to defend and protect the human body. In order to patrol the entire body, leukocytes travel by way of the circulatory system. The following cells are leukocytes of the innate immune system:



- *Phagocytes*, or *Phagocytic cells*: Phagocyte means “eating cell”, which describes what role phagocytes play in the immune response. Phagocytes circulate throughout the body, looking for potential threats, like bacteria and viruses, to engulf and destroy. You can think of phagocytes as security guards on patrol.
- *Macrophages*: Macrophages, commonly abbreviated as “Mφ”, are efficient phagocytic cells that can leave the circulatory system by moving across the walls of capillary vessels. The ability to roam outside of the circulatory system is important, because it allows macrophages to hunt pathogens with less limits. Macrophages can also release cytokines in order to signal and recruit other cells to an area with pathogens.



- **Mast cells:** Mast cells are found in mucous membranes and connective tissues, and are important for wound healing and defense against pathogens via the inflammatory response. When mast cells are activated, they release cytokines and granules that contain chemical molecules to create an *inflammatory cascade*. Mediators, such as histamine, cause blood vessels to dilate, increasing blood flow and cell trafficking to the area of infection. The cytokines released during this process act as a messenger service, alerting other immune cells, like neutrophils and macrophages, to make their way to the area of infection, or to be on alert for circulating threats.
- **Neutrophils:** Neutrophils are phagocytic cells that are also classified as *granulocytes* because they contain granules in their cytoplasm. These granules are very toxic to bacteria and fungi, and cause them to stop proliferating or die on contact.



The bone marrow of an average healthy adult makes approximately 100 billion new neutrophils per day. Neutrophils are typically the first cells to arrive at the site of an infection because there are so many of them in circulation at any given time.

- **Eosinophils:** Eosinophils are granulocytes target multicellular parasites. Eosinophils secrete a range of highly toxic proteins and free radicals that kill bacteria and parasites. The use of toxic proteins and free radicals also causes tissue damage during allergic reactions, so activation and toxin release by eosinophils is highly regulated to prevent any unnecessary tissue damage.

While eosinophils only make up 1-6% of the white blood cells, they are found in many locations, including the thymus, lower gastrointestinal tract, ovaries, uterus, spleen, and lymph nodes.

- **Basophils:** Basophils are also granulocytes that attack multicellular parasites. Basophils release histamine, much like mast cells. The use of histamine makes basophils and mast cells key players in mounting an allergic response.

- *Natural Killer cells*: Natural Killer cells (NK cells), do not attack pathogens directly. Instead, natural killer cells destroy infected host cells in order to stop the spread of an infection. Infected or compromised host cells can signal natural kill cells for destruction through the expression of specific receptors and antigen presentation.
- *Dendritic cells*: Dendritic cells are antigen-presenting cells that are located in tissues, and can contact external environments through the skin, the inner mucosal lining of the nose, lungs, stomach, and intestines. Since dendritic cells are located in tissues that are common points for initial infection, they can identify threats and act as messengers for the rest of the immune system by antigen presentation. Dendritic cells also act as bridge between the innate immune system and the adaptive immune system.

The Complement System

The complement system (also called the *complement cascade*) is a mechanism that *complements* other aspects of the immune response. Typically, the complement system acts as a part of the innate immune system, but it can work with the adaptive immune system if necessary. The complement system is made of a variety of proteins that, when inactive, circulate in the blood. When activated, these proteins come together to initiate the complement cascade, which starts the following steps:

1. *Opsonization*: Opsonization is a process in which foreign particles are marked for phagocytosis. All of the pathways require an antigen to signal that there is a threat present. Opsonization tags infected cells and identifies circulating pathogens expressing the same antigens.
2. *Chemotaxis*: Chemotaxis is the attraction and movement of macrophages to a chemical signal. Chemotaxis uses cytokines and chemokines to attract macrophages and neutrophils to the site of infection, ensuring that pathogens in the area will be destroyed. By bringing immune cells to an area with identified pathogens, it improves the likelihood that the threats will be destroyed and the infection will be treated.
3. *Cell Lysis*: Lysis is the breaking down or destruction of the membrane of a cell. The proteins of the complement system puncture the membranes of foreign cells, destroying the integrity of the pathogen. Destroying the membrane of foreign cells or pathogens weakens their ability to proliferate, and helps to stop the spread of infection.
4. *Agglutination*: Agglutination uses antibodies to cluster and bind pathogens together, much like a cowboy rounds up his cattle. By bringing as many pathogens together in the same area, the cells of the immune system can mount an attack and weaken the infection. Other innate immune system cells continue to circulate throughout the body in order to track down any other pathogens that have not been clustered and bound for destruction.

The steps of the complement cascade facilitate the search for and removal of antigens by placing them in large clumps, making it easier for other aspects of the immune system to do their jobs. Remember that the complement system is a supplemental cascade of proteins that assists, or “complements” the other aspects of the innate immune system.

The innate immune system works to fight off pathogens before they can start an active infection. For some cases, the innate immune response is not enough, or the pathogen is able to exploit the innate immune response for a way into the host cells. In such situations, the innate immune system works with the adaptive immune system to reduce the severity of infection, and to fight off any additional invaders while the adaptive immune system is busy destroying the initial infection.